

***Natural Remedies for Migraine Help**

***If you would like to have a complementary consultation with a natural health care provider in your area, to relieve your migraine headaches, please email info@release-the-pain.com .**

This report was written by Cathy Wong, ND, a specialist with over 14 years of research, clinical, and teaching experience in the field of alternative medicine. She is a corporate wellness consultant and you can also find this report in About.com.

What are Migraines?

According to the National Headache Foundation, more than 29.5 million Americans have migraines. Symptoms of migraine may include:

- Moderate to severe pain on one or both sides of the head
- Pulsating or throbbing pain
- Pain worse with physical activity
- Nausea with or without vomiting
- Sensitivity to light or sound

Approximately 20 percent of people with migraines experience what's called an aura prior to the migraine. Symptoms of an aura include flashes of light, zig zags, or blind spots in your vision or tingling in one arm or leg.

Natural Remedies for Migraine

1) Feverfew

The herb feverfew (*Tanacetum parthenium*) has been used for centuries as a folk remedy for headache, migraine, arthritis, pain, and fever in Europe. It became especially popular in England in the 1980s as an alternative to conventional drugs for migraines.

Several studies have evaluated the effectiveness of feverfew for the prevention and treatment of migraines. For example, one study examined the use of a feverfew extract or a placebo in 170 people with migraines. There was a significant decrease in migraine frequency with feverfew, which decreased by 1.9 migraines per month, compared to the placebo, which decreased by 1.3 migraines per month.

Not all studies have shown that feverfew is effective, however. A critical review of five studies on feverfew and migraines, involving a total of 343 people, concluded that results were mixed and did not convincingly establish that feverfew was effective for preventing migraine.

Side effects of feverfew may include abdominal pain, gas, diarrhea, nausea, vomiting, and nervousness. People with allergies to chamomile, ragweed, or yarrow should not take feverfew. Feverfew may increase bleeding time, so it should not be taken by people with bleeding disorders or two to three

weeks before or after surgery. The safety of feverfew in pregnant or nursing women or children has not been established.

Feverfew may interact with "blood-thinning" medications, such as aspirin and warfarin (Coumadin) and non-steroidal anti-inflammatory drugs (NSAIDs). People taking these medications should consult a healthcare practitioner before taking feverfew.

2) 5-HTP

5-HTP (5-Hydroxytryptophan) is a compound produced in the body from the amino acid tryptophan. It is used in the body to make the neurotransmitter serotonin and the hormone melatonin.

5-HTP is also available in supplement form. It is made from the seeds of an African plant, called *Griffonia simplicifolia*. Since supplements that affect levels of serotonin are used for migraines, 5-HTP has been explored as an alternative.

Some research indicates that 5-HTP may prevent migraines and reduce the frequency and severity of migraines, however large randomized controlled trials are needed. In one study, 124 people were given 5-HTP (600 mg/day) or the drug methysergide. After six months, 5-HTP was found to be as effective as methysergide in reducing the severity and duration of migraines

Another study looked at 5-HTP or the drug propranolol for 4 months. Both treatments resulted in a statistically significant reduction in the frequency of migraines. However, the propranolol group fared better, with a reduction in the duration of episodes and the number of analgesics used for the treatment of episodes. For more information on 5-HTP, read the 5-HTP Fact Sheet.

3) Magnesium

Magnesium is a mineral found naturally in foods such as green leafy vegetables, nuts, seeds, and whole grains and in nutritional supplements.

Magnesium is needed for more than 300 biochemical reactions. It helps regulate blood sugar levels and is needed for normal muscle and nerve function, heart rhythm, immune function, blood pressure, and for bone health.

Several studies have evaluated the effectiveness of magnesium for migraine and have had promising results. In one study, oral magnesium (600 mg per day) or a placebo were given to 81 people with migraines. After nine weeks, the frequency of migraines was reduced by 41.6 percent in the group taking magnesium, compared to 15.8 percent in the group taking the placebo. The only reported side effects were diarrhea in 18.6 percent of people and digestive irritation in 4.7 percent.

Other studies have found that magnesium reduces the severity and frequency of migraine. Not all studies have confirmed it, however, so more studies are needed. High doses of magnesium may cause diarrhea, nausea, loss of appetite, muscle weakness, difficulty breathing, low blood pressure, irregular

heart rate, and confusion. It can interact with certain medications, such as those for osteoporosis, high blood pressure (calcium channel blockers), as well as some antibiotics, muscle relaxants, and diuretics.

4) *Butterbur*

The herb butterbur (*Petasites hybridus*) is a shrub-like plant that grows in northern Asia, Europe, and parts of North America. Extracts made from the herb have been used to treat migraines, stomach cramps, coughs, allergies, and asthma.

Several studies suggest that butterbur helps to prevent migraines. The largest study involved 245 people with migraine took either butterbur extract (50 or 75 milligrams twice a day) or a placebo. After four months of treatment, butterbur, 75 milligrams twice a day but not 50 milligrams twice a day, was more effective than placebo for migraines. Side effects were mild and included mild digestive complaints, predominantly burping.

Side effects of butterbur may include indigestion, headache, fatigue, nausea, vomiting, diarrhea, or constipation. Pregnant or nursing women, children, or people with kidney or liver disease should not take butterbur.

Butterbur is in the ragweed plant family, so people who are allergic to ragweed, marigold, daisy, or chrysanthemum should not use butterbur. The raw herb as well as teas, extracts, and capsules made from the raw herb should not be used because they contain substances called pyrrolizidine alkaloids that can be toxic to the liver and kidneys and may cause cancer.

It is possible to remove the pyrrolizidine alkaloids from butterbur products. For example, in Germany, there is a safety limit to the level of pyrrolizidine alkaloids allowed in butterbur products. The daily recommended dose cannot exceed one microgram per day.

Sources:

Diener HC, Pfaffenrath V, Schnitker J, Friede M, Henneicke-von Zepelin HH. Efficacy and safety of 6.25 mg t.i.d. feverfew CO₂-extract (MIG-99) in migraine prevention--a randomized, double-blind, multicentre, placebo-controlled study. *Cephalalgia*. 25.11 (2005): 1031-1041.

Lipton RB, Gobel H, Einhaupl KM, Wilks K, Mauskop A. *Petasites hybridus* root (butterbur) is an effective preventive treatment for migraine. *Neurology*. 63.12 (2004): 2240-2244.

Peikert A, Wilimzig C, Kohne-Volland R. Prophylaxis of migraine with oral magnesium: results from a prospective, multi-center, placebo-controlled and double-blind randomized study. *Cephalalgia*. 16.4 (1996): 257-263.

Pfaffenrath V, Diener HC, Fischer M, Friede M, Henneicke-von Zepelin HH; Investigators. The efficacy and safety of *Tanacetum parthenium* (feverfew) in migraine prophylaxis--a double-blind, multicentre, randomized placebo-controlled dose-response study. *Cephalalgia*. 22.7 (2002): 523-532.

Pfaffenrath V, Wessely P, Meyer C, Isler HR, Evers S, Grotemeyer KH, Taneri Z, Soyka D, Gobel H, Fischer M. Magnesium in the prophylaxis of migraine--a double-blind placebo-controlled study. *Cephalalgia*. 16.6 (1996): 436-440.

Pittler MH, Ernst E. Feverfew for preventing migraine. *Cochrane Database Syst Rev*. 2004;(1):CD002286.

Wang F, Van Den Eeden SK, Ackerson LM, Salk SE, Reince RH, Elin RJ. Oral magnesium oxide prophylaxis of frequent migrainous headache in children: a randomized, double-blind, placebo-controlled trial. *Headache*. 43.6 (2003): 601-610.

DISCLAIMER:

The information contained on this report is for educational purposes only and should not be used as a substitute for diagnosis or treatment rendered by a licensed physician. It is essential that you discuss with your doctor any symptoms or medical problems that you may be experiencing.